



Bioinformatics Summer School

Long-reads Transcriptomics

Carolina Monzó

I2SysBio, Valencia, Spain

- 1 Introduction to Long-read Technologies
- 2 Experimental design
- 3 Use cases and wrap up


4

Section 1


Introduction to Long-read Technologies

Understanding the fundamentals of third-generation sequencing

illumina[®]



illumina[®]




Advantages:

- Higher accuracy
- Higher throughput
- Cost-effective

Disadvantages:

- Repetitive sequences
- Structural variants
- Isoform resolution – Alternative splicing



PacBio




Advantages:

- Single-molecule
- Isoform resolution – Alternative splicing
- Assembly simplification


Disadvantages:

- Lower accuracy
- Lower throughput
- Higher costs
- Bioinformatic challenges


PacBio





PacBio



Oxford NANOPORE Technologies









Section 2

Experimental design

Understanding the fundamentals of third-generation sequencing



WARNING!! Combination of sequencing platform and library preparation find **different transcripts**



WARNING!! Combination of sequencing platform and library preparation find **different transcripts**

What is your biological question?

Degraded RNA:



- Incomplete transcripts (difficulty to differentiate TSS and TTS)
- Biased coverage, tendency to over-represent 3' ends
- Compromised measurements of polyA tail lengths


RNA Integrity Number: ratio of intact ribosomal RNA peaks to degraded RNA fragments

- Get RIN as high as possible
- RIN > 7 is the minimum, but **RIN > 8 recommended** by ONT and PacBio
- Validate TTS and TSS with orthogonal data (CAGE-Seq or Quant-Seq)
- Incorporate RIN as a covariate in the Differential Expression analysis



Checking the RNA quality is not only important to get good quality full-length transcripts, but also to see the distribution of transcripts that we have in the transcriptome of our sample






Tissue extracted RNA integrity is not affected by the storage conditions in 14 days, and not affected by freeze-thaws up to 10 cycles.

Extract the RNA from your samples quickly!

Don't store the whole tissue, store the RNA



The RNA concentration increased over time at both room temperature and 4°C, showing that evaporation occurred at room temperature from day 3 and at 4°C from day 7.

Similarities between PacBio and ONT:

- Throughput (high number of reads per run)
- Capacity to sequence full-length transcripts

Technology	Platform	Median read length (kb)	Median throughput per run (10^6 transcript reads)
PacBio	cDNA+IsoSeq+Sequel II	~2.1	~2.6
	cDNA+Kinnex+Sequel II	~1.7	~40
	cDNA+Kinnex+Revio	~1.7	~100
ONT	cDNA+MinION (R10.4)	~0.939	~20
	cDNA+PromethION (R9.4)	~1	~130
	dRNA+MinION (R9.4)	~0.8	~1.1
	dRNA+PromethION (R9.4)	~0.6	~20

PacBio

- **Higher read accuracy**
- Commercial protocols have broad read length distribution
- Can't sequence dRNA or direct modifications



- Can sequence cDNA or dRNA
- Can simultaneously sequence dRNA and epitranscriptomic modifications
- Specific protocols can sequence ultra-long reads. **But standard protocols are biased towards short reads**
- Lower read accuracy
- Difficulty differentiating modifications


Number of reads per sample:


- Higher depth discovers more novel transcripts
- More complex organisms require more sequencing depth

Recommendations (human):

- 10 million reads for 80% of known isoforms
- > 20 million reads for detection of rare isoforms


$$\text{Depth}_{\text{sample}} = \frac{\sum \text{Depth}_{\text{isoform}}}{N_{\text{expressed isoforms in sample}}}$$






Bimodal distribution, **most** transcripts are present in **all** samples **or** are **sample-specific**, few are found in two or more samples

Biological Replicates



Replication is necessary for statistical power and to reduce the inclusion of spurious transcripts in downstream analyses.


High expression correlation between replicates





bc = 4 different barcodes for library prep
cx = 12 different barcodes for cDNA synthesis


Possible to have 48 samples in one run






For cDNA!!


Possible to have 24 samples in one run





Synthetic RNA molecules that mimic: isoforms, abundance, and transcript length

SIRVs and sequins





- Check if all SIRVs in the mix were detected in my sample (read length or depth biases)
- Check that quantification of SIRVs in the mix is as expected



- Non-blue reads in **spike-ins** represent the technical **noise level of the technology**
- Non-blue reads in the **mouse brains** are a combination of **technical and biological noise**

Use cases and wrap up

What's the most important thing to take into account when:

- 1. Studying drought effect on polyA length in Arabidopsis over time**
- 2. Reconstructing the transcriptome of a non-model organism**
- 3. Benchmarking bioinformatic tools for transcript quantification**

**What is the most important aspect
in the experimental design to
answer your biological question?**

- **Figure out your biological question before deciding on sequencing platform and library preparation method**
- **Use good quality RNA**
- **Sequence with enough depth**
- **Use biological replicates in your experiment**
- **Use synthetic controls**

Thank You!



For more information about the LongTREC Summer School:

<https://longtrec.eu>